

Fourier Transform Raman (FTR) Spectroscopy of Some Energetic Materials and Propellant Formulations II

Nicholas F. Fell
Jeffrey M. Widder
Stephen V. Medlin
Jeffrey B. Morris
Rose A. Pesce-Rodriguez
Kevin L. McNesby



ARL-TR-843

August 1995

19951011 066

NOTICES

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute indorsement of any commercial product.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE August 1995 3. REPORT TYPE AND DATES COVERED Final, January—April 1995 4. TITLE AND SUBTITLE Fourier Transform Raman (FTR) Spectroscopy of Some Energetic Materials and	
Fourier Transform Paman (FTP) Spectroscopy of Some Energetic Materials and	
Propellant Formulations II PR: 1L161102AH43	
6. AUTHOR(S)	
Nicholas F. Fell, Jeffrey M. Widder, Stephen V. Medlin, Jeffrey B. Morris, Rose A. Pesce-Rodriguez, and Kevin L. McNesby	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	ZATION
REPORT NUMBER	
U.S. Army Research Laboratory	
ATTN: AMSRL-WT-PC ARL-TR-843	
Aberdeen Proving Ground, MD 21005-5066	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING	ORING
AGENCY REPORT NUM	
11. SUPPLEMENTARY NOTES	:
12a. DISTRIBUTION / AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE	
Approved for public release; distribution is unlimited.	
Approved for public release; distribution is unlimited.	
Approved for public release; distribution is unlimited.	
13. ABSTRACT (Maximum 200 words)	1 :0 1:
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm	
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r	egion from
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline	egion from components
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline in a propellant formulation is demonstrated.	region from components e ingredient
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline in a propellant formulation is demonstrated. 14. SUBJECT TERMS 15. NUMBER OF 21	region from components e ingredient
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline in a propellant formulation is demonstrated. 14. Subject terms	region from components ingredient
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline in a propellant formulation is demonstrated. 14. SUBJECT TERMS Raman spectroscopy, energetic materials 15. NUMBER OF 21 16. PRICE CODE	region from components ingredient
13. ABSTRACT (Maximum 200 words) Fourier transform Raman (FTR) spectroscopy employing near infrared (NIR) laser radiation at 9,394.5 cm ⁻¹ characterize neat energetic materials and several propellant formulations. Raman spectra are reported over the r 100 cm ⁻¹ to 3,000 cm ⁻¹ , relative to the Rayleigh line. The technique is extended to the study of crystalline of propellant formulations during heating. The utility of the technique in determining the principal crystalline in a propellant formulation is demonstrated. 14. Subject terms 15. Number of 21	region from components components ingredient

ACKNOWLEDGMENT

Support is acknowledged from the National Research Council Postdoctoral Associateship Program and the Strategic Environmental Research and Development Program (SERDP) of the U.S. Department of Defense (DOD). Thanks are expressed to Dr. Monte Chawla of Atlas Wireline Services, Inc., for providing the samples of HNS and of the Chinese propellant formulation.

Accesio	n For	
NTIS DTIC Unanno Justific	TAB ounced	<u> </u>
By	ution /	
А	vailabilit	y Codes
Dist	A vail a Spe	
A-1		

TABLE OF CONTENTS

		<u>Page</u>
	ACKNOWLEDGMENT	iii
	LIST OF FIGURES	vii
1.	INTRODUCTION	1
2.	EXPERIMENTAL	1
3.	RESULTS AND DISCUSSION	3
3.1 3.2	Neat Energetic Materials	3 6
4.	CONCLUSION	11
5.	REFERENCES	13
	DISTRIBUTION LIST	15

LIST OF FIGURES

Figure		Page
1.	Simple diagram of the experimental apparatus used in these experiments	2
2.	The FTR spectra of RDX and HMX	4
3.	The FTR spectra of PETN and NQ	5
4.	The FTR spectra of ADN and TNT	5
5.	The FTR spectra of CL-20 and TNAZ	7
6.	The FTR spectra of HNS and pDNPA	7
7.	The FTR spectra of DMNA and NC	8
8.	The FTR spectra of GAP and AP	8
9.	The FTR spectra of RDX, HMX, and XM39	9
10.	The FTR spectra of RDX, XM39, and a propellant formulation believed to have been manufactured in mainland China	10
11.	The FTR spectra of XM39 as it is heated from 297 K to 458 K	10

1. INTRODUCTION

Fourier transform Raman (FTR) spectroscopy using near infrared (NIR) laser light as the scattering source has been shown to be useful in the identification of crystalline components of propellant formulations (Akhaven 1991). The ease with which many principal components of propellant formulations may be identified is due to the generally good scattering characteristics of crystalline energetic materials combined with the poor scattering characteristics of most plasticizers and binders. Although using NIR laser light as the scattering source limits the signal intensity compared to that obtained using visible laser light scattering, the lack of fluorescence interference when working in the NIR spectral region often outweighs this disadvantage. If a disadvantage exists in trying to measure Raman spectra of propellant formulations using NIR laser scattering, it is that most colored formulations will absorb the laser light and combust. Fortunately, most crystalline neat energetic materials are not colored, so the technique can often yield excellent results.

Recently we published a short paper consisting of Raman spectra of several neat energetic materials and several propellant formulations (McNesby et al. 1994). It was our goal at the time to have that paper be the first in a series to aid in the assembly of a Raman spectral database of energetic materials and propellant formulations. Since that time, modifications to our instrument have enabled coverage of a broader spectral range, so we have remeasured the spectra of most of the neat energetic materials reported in the first paper, added several more spectra of neat energetic materials not measured, and also have included some new energetic formulations and techniques. We hope most readers find this report a useful continuation of our initial effort towards the characterization of energetic materials by FTR spectroscopy.

2. EXPERIMENTAL

The experimental apparatus has been described previously (McNesby et al. 1994). Briefly, the equipment employed in these experiments consists of a Bomem DA-8.02 Fourier transform spectrometer to which a Raman accessory has been added. Incident radiation is provided by a Quantronix Series 100 Nd:YAG laser. A simple sketch of the experimental apparatus is shown in Figure 1. Raman-shifted radiation is collected using a back-scattering geometry and is detected after filtering and interferometer modulation using a liquid nitrogen-cooled InGaAs detector. Recently the interference filters used to exclude the Rayleigh line at a wavelength of 1.06 µm (9,394.5 cm⁻¹) were replaced by holographic notch




Figure 1. Simple diagram of the experimental apparatus used in these experiments.

filters (Kaiser Optical). This enabled measurement of Raman transitions shifted from 100 to 3,000 cm⁻¹ with respect to the Rayleigh line (the previous spectral range of the instrument was from 400 to 3,000 cm⁻¹). Also, we are able to observe anti-Stokes lines shifted from the Rayleigh line by 150 to 600 cm⁻¹.

All spectra reported here were measured at 4 cm⁻¹ resolution using coaddition of 256 scans. Incident laser power was 400 mW. Scan time to collect each spectrum was 11 min. All spectra were measured using the back scattering arrangement shown in Figure 1. Neat samples of energetic materials were placed in 1-mm i.d. glass capillary tubes, and the Nd:YAG laser focused on the surface of the tube. Propellant formulations, when in the form of cylindrical grains, were placed in a temperature-variable holder whose position could be maintained from run to run. The Nd:YAG laser light was then focused on a flat surface of the grain created by cleaving the grain normal to its longitudinal axis. Powdered propellant formulations were placed in a 5-mm-diameter glass tube that was then placed in a sample spinner (Princeton Photonics) operated at 2,400 rpm to minimize heating effects. The Nd:YAG laser was then

focused on the surface of the spinning tube. No correction was made to any of the spectra to account for responsivity of the detector, interferometer, or filters used in the experiments. For the spectra of neat energetic materials reported here, the intense feature at zero Raman shift is due to Rayleigh scatter that was not fully attenuated by the holographic filters. The intense ringing about zero shift is a manifestation of the intensity of the Rayleigh scatter entering the interferometer and the boxcar apodization used in the Fourier transformation of the raw data (Griffith and de Haseth 1986). Samples of HNS and of the Chinese propellant formulation were provided by Atlas Wireline Services, Inc. All other propellant samples used in these experiments were obtained from in-house sources.

3. RESULTS AND DISCUSSION

3.1 Neat Energetic Materials. Table 1 shows a summary of the energetic materials whose FTR spectra are described in this report. Figure 2 shows the FTR spectra of the most common nitramine energetic material, cyclotrimethylenetrinitramine (RDX), and of the eight-membered ring homolog, cyclotetramethylenetetranitramine (HMX). There are several interesting features in these spectra. First, unlike our previously published spectra, we are now able to observe in each spectrum vibrational transitions to within 100 cm⁻¹ of the Rayleigh line. Vibrational transitions due to ring deformations in RDX and HMX, occurring at Raman shifts of from 100 to 400 cm⁻¹, were not measurable in our laboratory prior to installation of the holographic rejection filters. Crystal lattice modes (Goetz and Brill 1979) in nitramine crystals, which occur below 100 cm⁻¹, fall within the "notch" of the holographic filters so are not observable. In the FTR spectrum of HMX, an intense peak is observed that is centered at a Raman shift of 150 cm⁻¹. This peak is close to the intensity of the most intense ring stretch features that occur in HMX from 800 to 1,000 cm⁻¹.

Figure 3 shows the spectrum of the nitrate ester pentaerythritol tetranitrate (PETN), a common ingredient in some commercial explosives. Also shown is the FTR spectrum of nitroguanidine (NQ). The N-H stretch in nitroguanidine at a Raman shift of approximately 3,200 cm⁻¹ is not observed since it occurs beyond the range of the InGaAs detector. Figure 4 shows the spectrum of ammonium dinitramide (ADN), an oxidizer whose use is rapidly expanding within the propellant community, and of trinitrotoluene (TNT). In both of these spectra, features below 400 cm⁻¹ are among the most intense in each spectrum. For the spectrum of TNT, the slightly elevated baseline is caused by sample heating during measurement of the spectrum since neat TNT is light yellow in color and absorbs some of the incident Nd:YAG laser radiation.

Table 1. Energetic Materials Used in These Experiments

Energetic Material	Laser Power (mW)	Composition
RDX	400	neat
HMX	400	neat
PETN	400	neat
NQ	400	neat
ADN	400	neat
TNT	400	neat
CL-20	400	neat
TNAZ	400	neat
HNS	400	neat
pDNPA	400	neat
NC	400	neat
DMNA	400	neat
AP	400	neat
GAP	400	neat
XM39	400	76% RDX,
		24% binders and plasticizers




Figure 2. The FTR spectra of RDX and HMX.




Figure 3. The FTR spectra of PETN and NQ.





Figure 4. The FTR spectra of ADN and TNT.

Raman Shift (wavenumbers)

Figure 5 shows the spectra of beta-hexanitrohexaazaisowurtzitane (CL-20, previously reported using the alternate name b-HNTW), an energetic caged nitramine, and 1, 3, 3-trinitroazetidine (TNAZ), a recently developed energetic nitramine containing a germinal dinitro group. Figure 6 shows the spectrum of hexanitrostilbene (HNS), an industrial explosive used in high-temperature applications, and polydinitropropyladapate (pDNPA), an energetic polymeric binder used in some propellant formulations. Both spectra were measured using incident radiation at a power of 400 mW, although the signal-to-noise ratio in the spectrum of HNS is superior to that of the spectrum of pDNPA. These spectra illustrate the general observation that crystalline energetic materials are better Raman scatterers than polymeric energetic materials (McNesby et al. 1994).

Figure 7 shows the FTR spectra of dimethylnitramine (DMNA), a simple energetic nitramine, and nitrocellulose (NC). While both spectra were measured using the same incident laser power and using the same sampling technique (1-mm i.d. capillary tube), the signal-to-noise ratio in the spectrum of crystalline DMNA is superior to that of polymeric nitrocellulose. Both samples were in the form of white powders. Figure 8 shows the FTR spectra of glycidyl azide polymer (GAP), an energetic binder, and ammonium perchlorate (AP), an important oxidizer for many composite propellant formulations, and the only inorganic energetic material included in this study. Again, it may be seen from Figure 8 that the FTR spectrum of the crystalline material (AP) has a higher signal-to-noise ratio than the polymeric GAP.

3.2 Applications to Propellant Analysis. Figure 9 shows the FTR spectra of RDX, HMX, and XM39, a propellant formulation whose principal ingredient is RDX. The spectrum of XM39 is an illustration of the utility of FTR spectroscopy for energetic material characterization in general and for identification of nitramine-based explosives in particular. While the spectra of RDX and HMX in Figure 2 were measured using neat powdered samples in capillary tubes, the spectrum of XM39 was measured by scattering the Nd:YAG laser light off of the interior surface of a cleaved propellant grain. Even with this simple sampling technique, assignment of RDX as the principal crystalline energetic ingredient in XM39 is unambiguous. Shown in the spectrum of XM39 is evidence of HMX "impurity" in the formulation. HMX is a normal by-product of RDX synthesis in most domestically produced RDX. Also shown in the spectrum of XM39 is the increased intensity (relative to neat RDX) of the feature near 150 cm⁻¹. This increased intensity (see Figure 2) may be due to the small HMX "impurity" in domestically manufactured RDX.




Intensity

Intensity


Raman Shift (wavenumbers)

Figure 5. The FTR spectra of CL-20 and TNAZ.



Raman Shift (wavenumbers)

Figure 6. The FTR spectra of HNS and pDNPA.



Intensity

Intensity

Figure 7. The FTR spectra of DMNA and NC.




Figure 8. The FTR spectra of GAP and AP.

Raman Shift (wavenumbers)




Figure 9. The FTR spectra of RDX, HMX, and XM39. Arrows indicate features in the spectrum of XM39 due to slight HMX "impurities" in domestically manufactured RDX.

Figure 10 shows the FTR spectra of RDX, XM39, and a commercially available propellant formulation believed to be manufactured in mainland China. Since the Chinese propellant formulation had a thin graphite coating that would cause the sample to absorb some of the incident laser radiation, the spectrum was measured by scattering the incident Nd:YAG laser radiation off of the surface of a spinning (2,400 rpm) 5-mm i.d. glass tube containing the propellant formulation. From Figure 10, it is apparent that the principal crystalline energetic material in the Chinese propellant formulation is RDX. Also, the spectrum of the Chinese propellant formulation shows no evidence of HMX, suggesting that the RDX present in the Chinese propellant formulation was manufactured by a different process than domestically produced RDX.

Figure 11 shows a portion of the FTR spectrum of the propellant formulation XM39 as it is heated from room temperature to the onset of decomposition at 458 K (185° C). This spectrum was measured by scattering the Nd:YAG laser light off the surface of a slice of XM39 taken from the interior of a propellant grain. As shown previously, the principal crystalline energetic ingredient in XM39 is RDX.




Figure 10. The FTR spectra of RDX, XM39, and a propellant formulation believed to have been manufactured in mainland China. Arrows indicate the presence of HMX in XM39.




Figure 11. The FTR spectra of XM39 as it is heated from 297 K to 458 K.

Note the slight red shift of the feature near 150 cm⁻¹.

Raman Shift (wavenumbers)

For XM39, as temperature is increased, the scattering intensity decreases (as shown by the decreased signal to noise ratio in the spectra) and there is a slight red shift of the feature near 150 cm⁻¹. This red shift was most notable for features at low Raman frequencies, and was negligible at Raman frequencies above 500 cm⁻¹. As the sample temperature approaches the melting point of RDX (477 K), the Raman intensity rapidly diminishes, reflecting a loss of crystallinity in the principal energetic ingredient in the formulation.

4. CONCLUSION

It has been shown that FTR spectroscopy using NIR laser radiation as the scattering source can be a useful tool in characterizing many neat samples of energetic materials and of some propellant formulations. Raman spectra of some common energetic materials, as well as some newly formulated energetic materials, have been measured from 100 to 3,000 cm⁻¹. To our knowledge, Raman spectra measured to within 100 cm⁻¹ of the Rayleigh line have not been reported for most energetic materials. Perhaps the most interesting result from our ability to measure Raman spectra over a wider spectral range is the observation that the most intense peak in the HMX spectrum occurs near 150 cm⁻¹. Finally, we believe that for many samples of energetic materials containing crystalline principal ingredients, FTR spectroscopy using NIR laser scattering should be the preferred choice of investigators for forensic analysis when adequate sample is present. We are presently extending the technique to the investigation of energetic materials at elevated temperatures, in formulations, and in solutions.

5. REFERENCES

- Akhaven, J. "Analysis of High Explosive Samples by Fourier Transform Raman Spectroscopy." Spectrochimica Acta. Part A, vol. 47, p. 1247, 1991.
- Goetz, F., and T. B. Brill. "Laser Raman Spectra of alpha-, beta-, gamma-, and delta-Octahydro-1,3,5,7-tetrazocine and Their Temperature Dependence." <u>Journal of Physical Chemistry</u>. Vol. 83, p. 340, 1979.
- Griffith, P. R., and J. A. de Haseth. <u>Fourier Transform Infrared Spectroscopy</u>. New York: John Wiley & Sons, 1986.
- McNesby, K. L., J. E. Wolfe, J. B. Morris, and R. A. Pesce-Rodriguez. "Fourier Transform Raman Spectroscopy of Some Energetic Materials and Propellant Formulations." <u>Journal of Raman Spectroscopy</u>. Vol. 25, p. 75, 1994.

NO. OF

COPIES ORGANIZATION

- 2 ADMINISTRATOR
 ATTN DTIC DDA
 DEFENSE TECHNICAL INFO CTR
 CAMERON STATION
 ALEXANDRIA VA 22304-6145
- 1 DIRECTOR
 ATTN AMSRL OP SD TA
 US ARMY RESEARCH LAB
 2800 POWDER MILL RD
 ADELPHI MD 20783-1145
- 3 DIRECTOR
 ATTN AMSRL OP SD TL
 US ARMY RESEARCH LAB
 2800 POWDER MILL RD
 ADELPHI MD 20783-1145
- 1 DIRECTOR
 ATTN AMSRL OP SD TP
 US ARMY RESEARCH LAB
 2800 POWDER MILL RD
 ADELPHI MD 20783-1145

ABERDEEN PROVING GROUND

5 DIR USARL ATTN AMSRL OP AP L (305)

NO. OF NO. OF COPIES ORGANIZATION COPIES ORGANIZATION COMMANDER HQDA ATTN AMSMI RD PR E A R MAYKUT ATTN SARD TT DR F MILTON AMSMI RD PR P R BETTS PENTAGON US ARMY MISSILE COMMAND WASHINGTON DC 20310-0103 REDSTONE ARSENAL AL 1 HODA ATTN SARD TT MR J APPEL OFFICE OF NAVAL RESEARCH DEPARTMENT OF THE NAVY PENTAGON ATTN R S MILLER CODE 432 WASHINGTON DC 20310-0103 800 N OUINCY STREET **ARLINGTON VA 22217** HODA OASA RDA ATTN DR C H CHURCH **COMMANDER** PENTAGON ROOM 3E486 ATTN J RAMNARACE AIR 54111C WASHINGTON DC 20310-0103 NAVAL AIR SYSTEMS COMMAND **WASHINGTON DC 20360** 4 COMMANDER ATTN R GHIRARDELLI **COMMANDER** D MANN ATTN R BERNECKER R 13 R SINGLETON G B WILMOT R 16 R SHAW NAVAL SURFACE WARFARE CENTER US ARMY RESEARCH OFFICE SILVER SPRING MD 20903-5000 P O BOX 12211 RSCH TRNGLE PK NC 27709-2211 COMMANDER 5 ATTN M C LIN 1 DIRECTOR J MCDONALD ATTN AMXRO MCS K CLARK E ORAN ARMY RESEARCH OFFICE J SHNUR P O BOX 12211 R J DOYLE CODE 6110 RSCH TRNGLE PK NC 27709-2211 NAVAL RESEARCH LABORATORY **WASHINGTON DC 20375** DIRECTOR ATTN AMXRO RT IP LIB SERVICES **COMMANDER** ARMY RESEARCH OFFICE ATTN T BOGGS CODE 388 P O BOX 12211 T PARR CODE 3895 RSCH TRNGLE PK NC 27709-2211 NAVAL WEAPONS CENTER CHINA LAKE CA 93555-6001 2 COMMANDER ATTN SMCAR AEE B D S DOWNS SUPERINTENDENT US ARMY ARDEC ATTN D W NETZER **PCTNY ARSNL NJ 07806-5000 DEPT OF AERONAUTICS** NAVAL POSTGRADUATE SCHOOL **COMMANDER MONTEREY CA 93940** ATTN SMCAR AEE J A LANNON US ARMY ARDEC PCTNY ARSNL NJ 07806-5000 AL LSCF ATTN R CORLEY R GEISLER COMMANDER 1 J LEVINE ATTN SMCAR AEE BR L HARRIS EDWARDS AFB CA 93523-5000 US ARMY ARDEC

PCTNY ARSNL NJ 07806-5000

NO. OF COPIES	ORGANIZATION	NO. OF COPIES	ORGANIZATION
1	AFOSR ATTN J M TISHKOFF BOLLING AIR FORCE BASE WASHINGTON DC 20332	2	PRINCETON COMBUSTION RESEARCH LABORATORIES INC ATTN N A MESSINA M SUMMERFIELD PRINCETON CORPORATE PLAZA
1	OSD SDIO IST ATTN L CAVENY PENTAGON WASHINGTON DC 20301-7100		BLDG IV SUITE 119 11 DEERPARK DRIVE MONMOUTH JUNCTION NJ 08852
1	COMMANDANT ATTN ATSF TSM CN USAFAS FORT SILL OK 73503-5600	3	DIRECTOR DIVISION 8354 ATTN S JOHNSTON P MATTERN D STEPHENSON SANDIA NATIONAL LABORATORIES
1	UNIV OF DAYTON RSCH INSTITUTE ATTN D CAMPBELL AL PAP	1	LIVERMORE CA 94550 BRIGHAM YOUNG UNIVERSITY
1	EDWARDS AFB CA 93523 NASA	1	ATTN M W BECKSTEAD DEPT OF CHEMICAL ENGINEERING PROVO UT 84058
	ATTN G B NORTHAM MS 168 LANGLEY RESEARCH CENTER LANGLEY STATION HAMPTON VA 23365	1	CALIFORNIA INSTITUTE OF TECH JET PROPULSION LABORATORY ATTN L STRAND MS 125 224 4800 OAK GROVE DRIVE
4	NATIONAL BUREAU OF STANDARDS ATTN J HASTIE		PASADENA CA 91109
	M JACOX T KASHIWAGI H SEMERJIAN US DEPARTMENT OF COMMERCE WASHINGTON DC 20234	1	CALIFORNIA INSTITUTE OF TECHNOLOGY ATTN F E C CULICK MC 301 46 204 KARMAN LAB PASADENA CA 91125
2	DIRECTOR ATTN C WESTBROOK W TAO MS L 282 LAWRENCE LIVERMORE NATIONAL LAB	1	UNIVERSITY OF CALIFORNIA LOS ALAMOS SCIENTIFIC LAB P O BOX 1663 MAIL STOP B216 LOS ALAMOS NM 87545
	P O BOX 808 LIVERMORE CA 94550	1	UNIVERSITY OF CALIFORNIA BERKELEY CHEMISTRY DEPARMENT ATTN C BRADLEY MOORE
	DIRECTOR ATTN B NICHOLS T7 MS B284 LOS ALAMOS NATIONAL LAB P O BOX 1663	1	211 LEWIS HALL BERKELEY CA 94720 UNIVERSITY OF CALIFORNIA SAN DIEGO
	LOS ALAMOS NM 87545	1	ATTN F A WILLIAMS AMES B010
_	NATIONAL SCIENCE FOUNDATION ATTN A B HARVEY WASHINGTON DC 20550		LA JOLLA CA 92093

NO. OF COPIES	ORGANIZATION	NO. OF COPIES	ORGANIZATION
2	UNIV OF CALIFORNIA SANTA BARBARA QUANTUM INSTITUTE ATTN K SCHOFIELD M STEINBERG SANTA BARBARA CA 93106	1	UNIVERSITY OF MICHIGAN GAS DYNAMICS LAB ATTN G M FAETH AEROSPACE ENGINEERING BLDG ANN ARBOR MI 48109-2140
	UNIV OF COLORADO AT BOULDER ENGINEERING CENTER ATTN J DAILY CAMPUS BOX 427 BOULDER CO 80309-0427	1	UNIVERSITY OF MINNESOTA DEPT OF MECHANICAL ENGINEERING ATTN E FLETCHER MINNEAPOLIS MN 55455
	UNIV OF SOUTHERN CALIFORNIA DEPT OF CHEMISTRY ATTN R BEAUDET S BENSON C WITTIG LOS ANGELES CA 90007	4	PENNSYLVANIA STATE UNIVERSITY DEPT OF MECHANICAL ENGINEERING ATTN K KUO M MICCI S THYNELL V YANG UNIVERSITY PARK PA 16802
_	CORNELL UNIVERSITY DEPARTMENT OF CHEMISTRY ATTN T A COOL BAKER LABORATORY ITHACA NY 14853	1	POLYTECHNIC INSTITUTE OF NY GRADUATE CENTER ATTN S LEDERMAN ROUTE 110 FARMINGDALE NY 11735
	UNIVERSITY OF DELAWARE ATTN T BRILL CHEMISTRY DEPARTMENT NEWARK DE 19711		PRINCETON UNIVERSITY FORRESTAL CAMPUS LIBRARY ATTN K BREZINSKY I GLASSMAN P O BOX 710
-	UNIVERSITY OF FLORIDA ATTN J WINEFORDNER DEPT OF CHEMISTRY GAINESVILLE FL 32611		PRINCETON NJ 08540 PURDUE UNIVERSITY SCHOOL OF AERONAUTICS AND ASTRONAUTICS
	GEORGIA INSTITUTE OF TECHNOLOGY SCHOOL OF AEROSPACE ENGINEERING ATTN E PRICE W C STRAHLE B T ZINN ATLANTA GA 30332	1	ATTN J R OSBORN GRISSOM HALL WEST LAFAYETTE IN 47906 PURDUE UNIVERSITY DEPARTMENT OF CHEMISTRY ATTN E GRANT WEST LAFAYETTE IN 47906
-	UNIVERSITY OF ILLINOIS DEPT OF MECH ENG ATTN H KRIER 144MEB 1206 W GREEN ST URBANA IL 61801	2	PURDUE UNIVERSITY SCHOOL OF MECHANICAL ENGINEERING ATTN N M LAURENDEAU S N B MURTHY TSPC CHAFFEE HALL
	THE JOHNS HOPKINS UNIV CPIA ATTN T W CHRISTIAN 10630 LITTLE PATUXENT PKWY SUITE 202		WEST LAFAYETTE IN 47906

COLUMBIA MD 21044-3200

NO. OF COPIES	ORGANIZATION	NO. OF COPIES	ORGANIZATION
1	RENSSELAER POLYTECHNIC INST DEPT OF CHEMICAL ENGINEERING ATTN A FONTUN TROY NY 12181	1	EXXON RESEARCH & ENG CO ATTN A DEAN ROUTE 22E ANNANDALE NJ 08801
1	STANFORD UNIVERSITY DEPT OF MECHANICAL ENGINEERING ATTN R HANSON STANFORD CA 94305	1	GENERAL APPLIED SCIENCE LABS INC 77 RAYNOR AVENUE RONKONKAMA NY 11779-6649
1	UNIVERSITY OF TEXAS DEPT OF CHEMISTRY ATTN W GARDINER AUSTIN TX 78712	1	GENERAL ELECTRIC ORDNANCE SYSTEMS ATTN J MANDZY 100 PLASTICS AVENUE PITTSFIELD MA 01203
1	VA POLYTECH INST AND STATE UNIV ATTN J A SCHETZ BLACKSBURG VA 24061	1	GENERAL MOTORS RSCH LABS PHYSICAL CHEMISTRY DEPARTMENT ATTN T SLOANE WARREN MI 48090-9055
1	APPLIED COMBUSTION TECHNOLOGY INC ATTN A M VARNEY P O BOX 607885 ORLANDO FL 32860	2	HERCULES INC ATTN W B WALKUP E A YOUNT P O BOX 210 ROCKET CENTER WV 26726
2	APPLIED MECHANICS REVIEWS ASME ATTN R E WHITE & A B WENZEL 345 E 47TH STREET NEW YORK NY 10017	1	HERCULES INC ATTN R V CARTWRIGHT 100 HOWARD BLVD KENVIL NJ 07847
1	ATLANTIC RESEARCH CORP ATTN R H W WAESCHE 7511 WELLINGTON ROAD GAINESVILLE VA 22065	1	ALLIANT TECHSYSTEMS INC MARINE SYSTEMS GROUP ATTN D E BRODEN MS MN50 2000 600 2ND STREET NE HOPKINS MN 55343
1	TEXTRON DEFENSE SYSTEMS ATTN A PATRICK 2385 REVERE BEACH PARKWAY EVERETT MA 02149-5900	1	ALLIANT TECHSYSTEMS INC ATTN R E TOMPKINS MN 11 2720 600 SECOND ST NORTH
1	BATTELLE TWSTIAC 505 KING AVENUE COLUMBUS OH 43201-2693	1	HOPKINS MN 55343 IBM CORPORATION ATTN A C TAM
1	COHEN PROFESSIONAL SERVICES ATTN N S COHEN 141 CHANNING STREET		RESEARCH DIVISION 5600 COTTLE ROAD SAN JOSE CA 95193
	REDLANDS CA 92373		IIT RESEARCH INSTITUTE ATTN R F REMALY 10 WEST 35TH STREET CHICAGO IL 60616

NO. OF COPIES	ORGANIZATION	NO. OF COPIES	ORGANIZATION
1	LOCKHEED MISSILES & SPACE CO ATTN GEORGE LO 3251 HANOVER STREET DEPT 52 35 B204 2 PALO ALTO CA 94304	1	SVERDRUP TECHNOLOGY INC ATTN J DEUR 2001 AEROSPACE PARKWAY BROOK PARK OH 44142
	OLIN ORDNANCE ATTN V MCDONALD LIBRARY P O BOX 222 ST MARKS FL 32355-0222 PAUL GOUGH ASSOCIATES INC	3	THIOKOL CORPORATION ELKTON DIVISION ATTN R BIDDLE R WILLER TECH LIB P O BOX 241 ELKTON MD 21921
	ATTN P S GOUGH 1048 SOUTH STREET PORTSMOUTH NH 03801-5423	3	THIOKOL CORPORATION WASATCH DIVISION ATTN S J BENNETT
1	HUGHES AIRCRAFT COMPANY ATTN T E WARD 8433 FALLBROOK AVENUE	1	P O BOX 524 BRIGHAM CITY UT 84302 UNITED TECHNOLOGIES RSCH CENTER
	CANOGA PARK CA 91303 ROCKWELL INTERNATIONAL CORP	1	ATTN A C ECKBRETH EAST HARTFORD CT 06108
	ROCKETDYNE DIVISION ATTN J E FLANAGAN HB02 6633 CANOGA AVENUE CANOGA PARK CA 91304	1	UNITED TECHNOLOGIES CORP CHEMICAL SYSTEMS DIVISION ATTN R R MILLER P O BOX 49028
	SCIENCE APPLICATIONS INC ATTN R B EDELMAN 23146 CUMORAH CREST WOODLAND HILLS CA 91364	1	SAN JOSE CA 95161-9028 UNIVERSAL PROPULSION COMPANY ATTN H J MCSPADDEN
_	SRI INTERNATIONAL ATTN G SMITH		25401 NORTH CENTRAL AVENUE PHOENIX AZ 85027-7837
	D CROSLEY D GOLDEN 333 RAVENSWOOD AVENUE MENLO PARK CA 94025	1	VERITAY TECHNOLOGY INC ATTN E B FISHER 4845 MILLERSPORT HIGHWAY EAST AMHERST NY 14051-0305
_	STEVENS INSTITUTE OF TECH DAVIDSON LABORATORY ATTN R MCALEVY III HOBOKEN NJ 07030	1	FREEDMAN ASSOCIATES ATTN E FREEDMAN 2411 DIANA ROAD BALTIMORE MD 21209-1525
	SVERDRUP TECHNOLOGY INC LERC GROUP ATTN R J LOCKE MS SVR 2 2001 AEROSPACE PARKWAY BROOK PARK OH 44142		ALLIANT TECHSYSTEMS ATTN J BODE 600 SECOND ST NE HOPKINS MN 55343

NO. OF NO. OF COPIES ORGANIZATION COPIES ORGANIZATION ABERDEEN PROVING GROUND 1 **ALLIANT TECHSYSTEMS** ATTN C CANDLAND 600 SECOND ST NE 36 DIR USARL HOPKINS MN 55343 ATTN: AMSRL-WT-P, A HORST AMSRL-WT-PC, **ALLIANT TECHSYSTEMS** R A FIFER 1 ATTN L OSGOOD **G F ADAMS** 600 SECOND ST NE W R ANDERSON R A BEYER HOPKINS MN 55343 S W BUNTE C F CHABALOWSKI **ALLIANT TECHSYSTEMS** 1 K P MC-NEILL BOONSTOPPEL ATTN R BURETTA A COHEN 600 SECOND ST NE HOPKINS MN 55343 R CUMPTON R DANIEL **ALLIANT TECHSYSTEMS** D DEVYNCK 1 N F FELL ATTN R BECKER B E FORCH 600 SECOND ST NE HOPKINS MN 55343 J M HEIMERL A J KOTLAR **ALLIANT TECHSYSTEMS** M R MANAA 1 ATTN M SWENSON W F MCBRATNEY K L MCNESBY 600 SECOND ST NE S V MEDLIN HOPKINS MN 55343 M S MILLER A W MIZIOLEK 1 US ARMY BENET LABORATORY ATTN SAM SOPOK S H MODIANO SMCAR CCB B J B MORRIS WATERVLIET NY 12189 J E NEWBERRY S A NEWTON R A PESCE-RODRIGUEZ B M RICE R C SAUSA M A SCHROEDER J A VANDERHOFF M WENSING A WHREN J M WIDDER

C WILLIAMSON AMSRL-CI-CA, R PATEL

USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. ARL Report Number _	ARL-TR-843	Date of Report August 1995
2. Date Report Received		
_		on purpose, related project, or other area of interest for which the report
4. Specifically, how is t	he report being use	d? (Information source, design data, procedure, source of ideas, etc.)
	-	quantitative savings as far as man-hours or dollars saved, operating costs lease elaborate.
organization, technical cor	ntent, format, etc.)	should be changed to improve future reports? (Indicate changes to
	Organization	·
CURRENT	Name	
ADDRESS	Street or P.O. Bo	ox No.
	City, State, Zip	Code
7. If indicating a Change of Old or Incorrect address b		ss Correction, please provide the Current or Correct address above and the
	Organization	
OLD ADDRESS	Name	
ADDRESS	Street or P.O. Bo	ox No.
	City, State, Zip	Code

(Remove this sheet, fold as indicated, tape closed, and mail.)
(DO NOT STAPLE)

DEPARTMENT OF THE ARMY

OFFICIAL BUSINESS



FIRST CLASS PERMIT NO 0001,APG,MD

POSTAGE WILL BE PAID BY ADDRESSEE

DIRECTOR
U.S. ARMY RESEARCH LABORATORY
ATTN: AMSRL-WT-PC
ABERDEEN PROVING GROUND, MD 21005-5066

NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES